arXiv:1503.06019v3 [nucl-th] 28 Oct 2016

Evolution of spatial anisotropies in quark gluon plasma

Golam Sarwar and Jan-e Alam
Theoretical Physics Division, Variable Energy Cyclotron Centre,
1/AF, Bidhan Nagar, Kolkata - 700064, INDIA

Evolution of spatial anisotropy created in the Heavy Ion Collisions at Relativistic Energies (HICRE) has been studied. The microscopic evolution of the fluctuations has been examined within the ambit of Boltzmann Transport Equation (BTE) in an expanding background. The expansion of the background composed of Quark Gluon Plasma (QGP) is treated within the framework of relativistic hydrodynamics. Spatial anisotropies of the initial state with different geometry have been evolved through Boltzmann equation and studied how these anisotropies dissipates during their propagation. This is very useful in understanding the presumption that the measured anisotropies in the data imitate the initial state effects. Relation between thermal fluctuation and transport coefficients like, shear viscosity and thermal conductivity have been presented. We have also estimated the angular power spectrum of the anisotropies in the appendix.

PACS numbers: 25.75.-q,25.75.Dw,24.85.+p

I. INTRODUCTION

The main aim of the heavy ion collision experiments at Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC) is to create a new state of matter called quark gluon plasma (QGP). Such a state of matter, *i.e.* QGP may have existed in the early universe after a few micro-second of the big bang. One of the motivation to create and study QGP in the laboratory is to understand the state of the universe in the micro-second old era. The temperature fluctuation in the cosmic microwave background radiation (CMBR) has provided crucial information about the universe when it was about 300,000 years old. This information has led to tremendous support to the big bang model of the cosmology. The polarization of the photons resulting from the Thomson scattering at the decoupling surface infected by density fluctuation gets reflected in the quadrupole moment of the phase space distribution of the incident photon. The temperature fluctuation in the CMBR is introduced as a perturbation in the phase space distribution of photons. The evolution of this perturbation is studied by using Boltzmann transport equation (BTE) [1] in gravitational field with Thomson scattering in the collision term. The linear polarization resulted from the scattering is connected with the quadrupole moment of the photon's phase space distribution.

We have adopted a similar approach here to study the spatial anisotropy implemented by introducing a deviation, $\delta f(x,p,t)$ in the equilibrium distribution function, $f^{(0)}(p)$. The bulk in equilibrium evolves via relativistic hydrodynamics and the evolution of the fluctuations over the equilibrated system, δf is treated within the ambit of BTE. In contrast to this the propagation of perturbation has been studied using hydrodynamics in [2] (see also [3]). δf may be used to estimate the fluctuations in various thermodynamical quantities as we will see below. In the present work the equilibrated background is assumed to be quark gluon plasma (QGP) expected to be produced in heavy ion collision at relativistic energies (HICRE).

The fluctuations in the thermodynamic quantities (e.g. hot spots created in the initial state of the collisions [4]) can be related to perturbations in the phase space distributions in the hydrodynamic limit. The evolution of these fluctuations can be analyzed within the ambit of BTE [5]. Fluctuations in thermodynamic quantities have been proposed as signals of the critical end point in the QCD phase transition [6, 7]. Dissimilar fluctuations in partonic and hadronic phases in the net electric charge and baryon number may shed light on the QCD phase transition in HICRE [8]. Event by event fluctuations in the ratio of positively to negatively charged pions may be used as an indicator of QCD transition [9] as well as for understanding the chemical equilibrium in the system formed in HICRE [10]. Evolution of these fluctuations near the critical end point has been studied by using BTE [11]. Kinetic theory approach has also been adopted to study fluctuations in particle and energy densities [12]. In Ref. [13] it has been argued that the perturbations in hot QGP travel longer distance to reach the border of the medium giving rise to the possibility of detectable signatures of these perturbations.

The study of the fluctuations in the space time structure of the fireball driven by the fluctuations in the position of the nucleons in the colliding nuclei is an important contemporary issue in HICRE. Fluctuations in the space-time structure of the system will infect fluctuations in the thermodynamic quantities. How will these fluctuations evolve with time in a hydrodynamically expanding system and how are they connected with the transport coefficients are addressed in this work.

The paper is organized as follows. In the next section we discuss the evolution of fluctuations in a non-expanding background and present a relation between energy fluctuation in Fourier space and viscous coefficient. Section III is devoted to discuss the progression of fluctuations in a hydrodynamically expanding QGP background. Results are presented in section IV and section V is dedicated to summary and discussions.

II. EVOLUTION OF FLUCTUATIONS IN A NON-EXPANDING BACKGROUND

In the following subsections we discuss the connection of δf, deviation of phase space distribution from its equilibrium value with the fluctuations in various thermodynamic variables and its time evolution in a non-expanding background within the framework of BTE. The phase space distribution function, $f(\vec{x},\vec{p},t)$ of a system slightly away from equilibrium, at time t, position $\vec{x}$, momentum $\vec{p}$ can be written as [14],

$$f(\vec{x},\vec{p},t) = f^{(0)}(p)\{1+\Psi(\vec{x},\vec{p},t)\}, \tag{1}$$

where $f^{(0)}(p)$ is the equilibrium phase space distribution function determined by the local temperature, $T(t,\vec{x})$ and $\Psi(\vec{x},\vec{p},t)$ is the fractional deviation from $f^0(p)$. $\Psi(\vec{x},\vec{p},t)$ can be used to estimate the fluctuations in various thermodynamic quantities in the system. The evolution of Ψ is governed by BTE, which in turn provides the relation between the dissipative effects and fluctuations in the hydrodynamic limit.

A. Fluctuations in various hydrodynamic quantities in Fourier space

The energy momentum tensor, $T^{\mu\nu}$ of the system under study can be written as: $T^{\mu\nu} = \overline{T}^{\mu\nu} + \Delta T^{\mu\nu}$, where the equilibrium (ideal) part, $\overline{T}^{\mu\nu}$ is determined by $f^0(p)$ and the dissipative part, $\Delta T^{\mu\nu}$ is determined by Ψ, *i.e.*

$$T^{\mu\nu} = \int d^3p\, \frac{p^\mu p^\nu}{p^0} f(\vec{x},\vec{p},t) \tag{2}$$

where $f(\vec{x},\vec{p},t)$ is given by Eq. 1. The $\overline{T}^{\mu\nu}$ of the system in equilibrium can be obtained from Eq. 2 by setting $\Psi=0$. The metric, $g^{\mu\nu}$, in Minkowski space-time is taken as $g^{\mu\nu}=(-1,1,1,1)$. We assume that the momentum, $\vec{p}$ can be written as $p_i = pn_i$, *i.e.* $\vec{p}=|\vec{p}|\hat{n}$ where $\hat{n}$ is a unit vector and $d^3p = p^2 dp\, d\Omega$, $d\Omega$ being solid angle associated with n_i which satisfy $\int d\Omega n_i n_j = 4\pi\delta_{ij}/3$ and $\int d\Omega n_i n_j n_k = 0$.

It is straightforward to obtain various components of T^μ_ν from Eq. 2. The deviation of the components of the stress energy tensor, $\delta T^{\mu\nu}$ from their ideal values can be expressed in terms of the perturbation, Ψ as follows:

$$\begin{aligned}
\delta T^0_0 &= -\int p^2 dp\, d\Omega\, \epsilon\, f^{(0)}(p)\Psi(\vec{x},\vec{p},t), \\
\delta T^0_i &= \int p^2 dp\, d\Omega\, p n_i f^{(0)}(p)\Psi(\vec{x},\vec{p},t), \\
\delta T^i_j &= \int p^2 dp\, d\Omega\, \frac{p^2}{\epsilon}\, n^i n_j\, f^{(0)}(p)\Psi(\vec{x},\vec{p},t)
\end{aligned} \tag{3}$$

where $\epsilon = p^0 = -p_0 = \sqrt{p^2+m^2}$ and m is the mass of the particle. It is to be noted that the integral, $\int d^3p\, p n_i f^{(0)}(p) = 0$ since in equilibrium the system is isotopic i.e, all directions are equally probable in its rest frame, the phase space average of momentum vector then is zero.

The ideal part of the stress energy tensor in the hydrodynamic limit is given by,

$$\overline{T}^\mu_\nu = (\bar{\rho}+\bar{P})U^\mu U_\nu + \bar{P} g^\mu_\nu, \tag{4}$$

where $U^\mu = dx^\mu/d\tau$ is the four velocity of the fluid and τ is the proper time. The relations among various components of $\overline{T}^{\mu\nu}$ and the thermodynamic variables are: $\overline{T}^0_0 = -\bar{\rho}$, $\overline{T}^0_i = -\bar{T}^i_0 = 0$, and $\overline{T}^i_j = \bar{P}\delta^i_j$. Therefore, $\bar{\rho} = \int p^2 dp\, d\Omega\, \epsilon f^{(0)}(p)$, is the average energy density. Other thermodynamic quantities like pressure, number density etc can be estimated in a similar way. Equilibrium distribution is isotropic, therefore, integration over $d\Omega$ will give simply 4π. If the fluid is slightly away from equilibrium with space time dependent fluctuations in energy density, pressure, velocity, etc., then the system will evolve toward equilibrium through dissipative processes. In such situation the components of the energy momentum tensors can be explicitly expressed in terms of the thermodynamic variables as:

$$\begin{aligned} T^0_0(x_i,t) &= -\{\bar{\rho}+\delta\rho(x_i,t)\}, \\ T^0_i(x_i,t) &= -T^i_0 = (\bar{\rho}+\bar{P})v_i, \\ T^i_j(x_i,t) &= \{\bar{P}+\delta P(x_i,t)\}\delta^i_j + \Sigma^i_j(x_i,t), \\ \Delta T^i_i &= 0, \end{aligned} \tag{5}$$

where v_i is the i^{th} component of the velocity perturbation. We can choose a frame which is moving with velocity close to the velocity of the fluid, so that the fluid velocity measured from this frame is small. From Eq. 5, we get $\Sigma^i_j = T^i_j - \delta^i_j T^k_k/3$. By using Eqs. 3, 4 and 5 we get,

$$\begin{aligned} \delta\rho(x_i,t) &= -\delta T^0_0(x_i,t), \\ v_i(x_i,t) &= \frac{\delta T^0_i(x_i,t)}{(\bar{\rho}+\bar{P})}, \\ \delta T^i_j(x_i,t) &= \delta P(x_i,t)\delta^i_j + \Sigma^i_j(x_i,t), \end{aligned} \tag{6}$$

with $\delta P(x_i,t) = \delta T^i_i(x_i,t)/3$. The shear stress, $\Sigma^i_j(x_i,t)$ can be expressed in terms of shear viscous coefficient, η as $\Sigma^i_j(x_i,t) = -\eta(\frac{\partial U_i}{\partial x_j} + \frac{\partial U_j}{\partial x_i} - \frac{2}{3}\delta^i_j\frac{\partial U_l}{\partial x_l})$ and the thermal conductivity (χ) is defined through the relation, $\delta T^0_j = -\chi(\frac{\partial T}{\partial x_i} + T\frac{\partial U}{\partial t})$ [15]. It is useful to express these quantities, *i.e.* various components of $\delta T^{\mu\nu}$ in Fourier or k-space because expansion of these quantities in terms the spherical harmonics (Y_{lm}) will enable to connect the angular scales set by l in terms of k analogous to the determination of angular scale in CMBR [1]. In $k-$space these quantities are marked by $\tilde{}$ as: $\tilde{\Sigma}^l_j(k_i,t) = -i\eta(\tilde{U}^l k_j + \tilde{U}_j k^l - \frac{2}{3}\delta^l_j k^r\tilde{U}_r)$ and $\delta\tilde{T}^0_i(k_i,t) = -\chi\left[ik_i\tilde{T}(k_l,t) + \tilde{T}(k_l,t)\frac{\partial\tilde{U}_i}{\partial t}\right]$.

By using Eqs. 3 and 6 the fluctuations in momentum space can be expressed as:

$$\begin{aligned} \delta\tilde{\rho}(k_i,t) &= \int p^2 dp\, d\Omega\, \epsilon\, f^{(0)}(p)\tilde{\Psi}(k_i,p,n_i,t), \\ \tilde{v}_i(k_l,t) &= -\frac{1}{(\bar{\rho}+\bar{P})}\int p^2 dp\, d\Omega\, pn_i f^{(0)}(p)\tilde{\Psi}, \\ \tilde{\Sigma}^i_j(k_l,t) &= \int p^2 dp\, d\Omega\,\frac{p^2}{\epsilon}(n_in_j - \frac{1}{3}\delta_{ij})f^{(0)}(p)\tilde{\Psi}, \\ \delta\tilde{P}(k_i,t) &= \frac{1}{3}\int p^2 dp\, d\Omega\,\frac{p^2}{\epsilon}f^{(0)}(p)\tilde{\Psi} \end{aligned} \tag{7}$$

where $\tilde{\Psi}$ is the Fourier transform of Ψ. Now we take the zenith direction along $\vec{k}$ and then the angular dependence of $\tilde{\Psi}(k_i,p,n_i,t)$ can be expressed in terms of angles between $\hat{k}$ and $\vec{n}$. Depending on the symmetries of the problem under consideration $\tilde{\Psi}$ can be expanded in a series of suitably chosen basis functions e.g, for axial symmetry in terms of Legendre polynomials and in absence of such symmetry it can be expressed in terms of spherical harmonics [16].

The vector component n_i and tensor components $(n_in_j - \frac{1}{3}\delta_{ij})$, appearing in the expressions for $v_i(k_l,t)$ and $\Sigma^i_j(k_l,t)$ respectively, can be converted into functions of θ (angle between $\vec{k}$ and $\vec{n}$), by taking contraction with suitable tensors made out of the components of $\hat{k}$. If we contract n_i with k_i then we get $k\hat{k}\cdot\hat{n} = k\cos\theta = kP_1(\hat{k}\cdot\vec{n})$ and by contracting $(n_in_j - \frac{1}{3}\delta_{ij})$ with $(\hat{k_i}\hat{k_j} - \frac{1}{3}\delta_{ij})$ we get $\frac{2}{3}\frac{1}{2}(3(\hat{k}\cdot\hat{n})^2 - 1) = \frac{2}{3}\frac{1}{2}(3\cos^2\theta - 1) = \frac{2}{3}P_2(\hat{k}\cdot\hat{n})$, where P_l s are Legendre polynomials. For axial symmetric distribution of $\vec{p}$ it helps to connect different co-efficients of expansion of $\tilde{\Psi}$ in terms of Legendre Polynomials with corresponding scalar quantities obtained from $v_i(k_l,t)$ and $\Sigma^i_j(k_l,t)$ due to orthogonality relation satisfied by P_l's. We define the scalar quantities like Δ, θ and σ as [17],

$$\Delta(k_i,t) = \frac{\delta\tilde{\rho}(k_i,t)}{\bar{\rho}} = -\frac{\delta\tilde{T}^0_0(k_i,t)}{\bar{\rho}}, \tag{8}$$

is the fluctuation in energy density in Fourier space. Similarly we define energy flux,

$$\theta(k_i,t) = ik^j\tilde{v}_j = \frac{ik^j\delta\tilde{T}^0_j(k_i,t)}{(\bar{\rho}+\bar{P})}, \tag{9}$$

and the shear stress as,

$$(\bar{\rho}+\bar{P})\sigma(k_l,t) = -(\hat{k_i}\hat{k_j} - \frac{1}{3}\delta_{ij})\tilde{\Sigma}^i_j(k_l,t), \tag{10}$$

The quantity, $\theta(k_i,t) = ik^jv_j$ originates from the velocity gradient. θ and σ can be expressed in terms of the shear viscous coefficient (η) and thermal conductivity (χ) as follows:

$$\sigma(\vec{k},t) = -\frac{4}{3}\frac{\eta}{\bar{\rho}+\bar{P}}\, ik^j\tilde{U}_j(\vec{k},t) \tag{11}$$

and

$$\theta(\vec{k},t) = \frac{\chi}{\bar{\rho}+\bar{P}}(k^2\tilde{T}(k_l,t) - ik_l\tilde{T}(k_l,t)\dot{\tilde{U}}_l(k_l,t)). \tag{12}$$

Now Eqs. 7, 8, 9 and 10 can be used to obtain the fluctuations in the energy density, pressure and velocity in k-space as:

$$\begin{aligned}
\Delta(k_i,t) &= \frac{1}{4\pi}\int d\Omega\,\frac{\int p^2dp\,\epsilon f^{(0)}(p)\tilde{\Psi}(k_i,p,n_i,t)}{\int p^2dp\,\epsilon f^{(0)}(p)},\\
\frac{\delta\tilde{P}(k_i,t)}{\bar{P}} &= \frac{1}{4\pi}\int d\Omega\,\frac{\int p^2dp\,\frac{p^2}{\epsilon}f^{(0)}(p)\tilde{\Psi}(k_i,p,n_i,t)}{\int p^2dp\,p^2/\epsilon f^{(0)}(p)},\\
\theta(k_i,t) &= \frac{ik}{4\pi}\int d\Omega\,(\hat{k}.\hat{n})\,\frac{\int p^2dp\,f^{(0)}(p)\tilde{\Psi}(k_i,p,n_i,t)}{\int p^2dp\,(\epsilon+p^2/3\epsilon)f^{(0)}(p)},\\
\sigma(k_i,t) &= -\frac{1}{4\pi}\int d\Omega\,((\hat{k}.\hat{n})^2-\frac{1}{3})\frac{\int p^2dp\,f^{(0)}(p)\tilde{\Psi}(k_i,p,n_i,t)}{\int p^2dp\,(\epsilon+p^2/3\epsilon)f^{(0)}(p)}.
\end{aligned} \tag{13}$$

To understand the angular scale determined by the multipole number l (as used in the appendix to find the angular correlations) we expand $\tilde{\Psi}$ in terms of Legendre polynomials for an axially symmetric distribution as:

$$\tilde{\Psi}(\vec{k},\hat{n},p,t) = \sum_{l=0}^{\infty}(-i)^l(2l+1)\Psi_l(\vec{k},p,t)P_l(\hat{k}.\hat{n}), \tag{14}$$

where the factor$(-i)^l(2l+1)$ is used to simplify the expansion of a plane wave form of $\tilde{\Psi}$. The l is related to angular resolution of the anisotropies, *i.e.* smaller angular scale will require larger l and vice versa. The temperature fluctuations may be written in terms of Ψ as $\Delta T/\bar{T} = -(\partial lnf/\partial lnp)\Psi$ [17]. For simplicity we will consider the massless limit, $m=0$ which gives $\epsilon = p$. The energy density is a scalar quantity whereas the velocity and the shear tensor are vector and tensor respectively and these aspects are also bound to reflect in the corresponding fluctuations. Therefore, the orthogonality of P_l's ensure that the fluctuations in scalar, vector and tensor quantities are dictated by the coefficients Ψ_0, Ψ_1 and Ψ_2 which are obtained by substituting Ψ from Eq. 14 in Eq. 13 and performing the angular integration as,

$$\begin{aligned}
\Delta(k_i,t) = \delta\rho(k_i,t)/\bar{\rho} &= \int p^2dp\,pf^{(0)}(p)\Psi_0(k_i,p,t)/\bar{\rho},\\
\delta P(k_i,t)/\bar{P} &= \int p^2dp\,pf^{(0)}(p)\Psi_0(k_i,p,t)/\bar{\rho}\\
\theta(k_i,t) = 3ik^j\delta T^0_j/(4\bar{\rho}) &= \frac{3}{4}k\int p^2dp\,pf^{(0)}(p)\Psi_1(k_i,p,t)/\bar{\rho}\\
\sigma(k_l,t) &= \frac{1}{2}\int p^2dp\,pf^{(0)}(p)\Psi_2(k_i,p,t)/\bar{\rho}
\end{aligned} \tag{15}$$

where $\bar{\rho} = \int p^2 dp\, p f^{(0)}(p)$. The above set of equations can be written in a more compact form as:

$$F(\hat{n}, \vec{k}, t) = \int p^2 dp\, p f^{(0)}(p) \Psi(\vec{k}, p, n_i, t)/\bar{\rho} \tag{16}$$

therefore, F has the angular dependence of Ψ and consequently F can be expressed as:

$$F(\vec{k}, \hat{n}, t) = \sum_{l=0}^{\infty} (-i)^l (2l+1) F_l(\vec{k}, t) P_l(\hat{k}.\hat{n}). \tag{17}$$

with

$$F_l(\vec{k}, t) = \int p^2 dp\, p f^{(0)}(p) \Psi_l(\vec{k}, p, t)/\epsilon_0 \tag{18}$$

The fluctuations in terms of F_l's are now given by

$$\Delta(\vec{k}, t) = F_0(\vec{k}, t),\ \delta P(\vec{k}, t)/\bar{P} = F_0(\vec{k}, t),\ \theta(\vec{k}, t) = 3/4kF_1(\vec{k}, t),\ \sigma(\vec{k}, t) = 1/2F_2(\vec{k}, t). \tag{19}$$

Using the relation $\sigma(\vec{k}, t) = -4\eta i k^j U_j(\vec{k}, t)/3(\bar{\rho} + \bar{P})$, and writing $ik^j U_j(\vec{k}, t) = \Theta(\vec{k}, t)$ we get an important relation which connects the fluctuation (F_2) with the transport coefficient (η),

$$F_2(\vec{k}, t) = -\frac{8\eta}{3(\bar{\rho} + \bar{P})} \Theta(\vec{k}, t) \equiv -\frac{8}{3\bar{T}} \frac{\eta}{s} \Theta(\vec{k}, t) \tag{20}$$

where the thermodynamic relation, $\bar{\rho} + \bar{P} = sT$ has been used. In Eq.20 s is the entropy density and $\bar{T}$ is the average temperature. The η appears as a coefficient of 2^{nd} rank tensor involving gradient in the ith direction of the jth component of velocity, as a result the $l = 2$ term appears in the expression for η in 20.

B. Fluctuations in Fourier space and transport coefficients in relaxation time approximation

The temperature fluctuations, $\delta T(\theta, \phi)$ in CMBR is generally expanded in Laplace series in terms of spherical harmonics, $Y_{lm}(\theta, \phi)$. The maximum value of l is determined by the angular resolution of the detector which can be connected to the wave number (k) corresponding to the Fourier transform of the spatial anisotropy. Therefore, in analogy with fluctuations in the CMBR the spatial anisotropy is studied here in Fourier space. But first we briefly discuss it in coordinate space.

The BTE, $p^\mu \partial_\mu f = C[f]$ in absence of external force and in the relaxation time approximation (similar approximation were used *e.g.* in [18–20]) used reduces to

$$\frac{\partial \Psi}{\partial t} + \frac{p^i}{\epsilon} \frac{\partial \Psi}{\partial x^i} = -\frac{\Psi(\vec{x}, \vec{p}, t)}{\tau_R}. \tag{21}$$

for Ψ. In Eq.21 τ_R is the relaxation time. In this work we assume that the system is close to the local equilibrium and the collisions between the particles bring the system back to the equilibrium within a time scale τ_R. For the present work the relaxation time can be estimated as the inverse of the reaction rate of the quarks and gluons using pQCD cross sections and Hard Thermal Loop Approximations [21]. The solution of Eq. 21 for a given initial (at time t_0) distribution, $\Psi_{in}(\vec{x}, \vec{p}, t_0)$ is [22]:

$$\Psi(\vec{x}, \vec{p}, t - t_0) = \Psi_{in}\left((\vec{x} - \frac{\vec{p}}{\epsilon}(t - t_0)), \vec{p}\right) \exp\left[-\frac{(t - t_0)}{\tau_R}\right] \tag{22}$$

Knowing Ψ it is straightforward to estimate the fluctuation in energy density from the following expression:

$$\Delta(\vec{x}, t - t_0) = \frac{\int p^2 dp\, d\Omega\, \epsilon f^{(0)}(p) \Psi(\vec{x}, \vec{p}, t - t_0)}{\int p^2 dp\, d\Omega\, \epsilon f^{(0)}(p)}. \tag{23}$$

The solution 22 is useful to study the time evolution of spatial anisotropy of the matter.

Now we would like to derive a relation between the fluctuation in energy density and transport coefficients in a non-expanding system. To facilitate this we write Eq. 21 for massless particles (as the case may be for partonic plasma produced in HICRE) in momentum space:

$$\frac{\partial \Psi}{\partial t} + ik(\hat{k}.\hat{n})\Psi = -\frac{\Psi(\vec{k},\vec{p},t)}{\tau}. \tag{24}$$

With the help of Eq. 16, Eq. 24 can be reduced to an equation describing the time evolution of F as

$$\frac{\partial F}{\partial t} + ik(\hat{k}.\hat{n})F = -\frac{F(\vec{k},t)}{\tau}. \tag{25}$$

This equation has the following solution,

$$F(\vec{k},\hat{n},t) = F(\vec{k},\hat{n},t_0)\exp\left[-(\frac{1}{\tau}+ik\mu)(t-t_0)\right] \tag{26}$$

where $\hat{k}.\hat{n}=\mu$. The value of $F(\vec{k},\hat{n},t)$ can be obtained from its value at initial time, t_0. Eq. 26 is a general expression for the fluctuations in the sense that all the quantities, *e.g.* Δ, θ, σ, discussed above at time t can be obtained from this expression if their corresponding initial values are supplied. Expanding $F(\vec{k},\hat{n},t)$ as in Eq. 17 and using the orthogonality relations of $P_l(\mu)$s we get,

$$F_l(\vec{k},t) = \frac{1}{2}e^{-\frac{(t-t_0)}{\tau}}\sum_{s=0}^{\infty}(-i)^{(s-l)}(2s+1)F_s(\vec{k},t_0)\int_{-1}^{+1}d\mu P_l(\mu)P_s(\mu)e^{-ik\mu(t-t_0)} \tag{27}$$

Therefore, the energy density fluctuation at time t is given by:

$$\Delta(\vec{k},t) = \frac{1}{2}e^{-\frac{(t-t_0)}{\tau}}\sum_{s=0}^{\infty}(-i)^{s}(2s+1)F_s(\vec{k},t_0)\int_{-1}^{+1}d\mu P_s(\mu)e^{-ik\mu(t-t_0)}. \tag{28}$$

Taking terms upto $s=2$ in the expression for $\Delta(\vec{k},t)$ we get,

$$\Delta(\vec{k},t) = \frac{1}{2}e^{-\frac{(t-t_0)}{\tau}}\sum_{s=0}^{2}(-i)^{s}(2s+1)F_s(\vec{k},t_0)\int_{-1}^{+1}d\mu P_s(\mu)e^{-ik\mu(t-t_0)}. \tag{29}$$

Performing the integration over μ we get,

$$\begin{aligned}\Delta(\vec{k},t) =& e^{-\frac{(t-t_0)}{\tau}}[F_0(\vec{k},t_0)\{\frac{\sin k(t-t_0)}{k(t-t_0)}\} + 3F_1(\vec{k},t_0)\{\frac{\cos k(t-t_0)}{k(t-t_0)} - \frac{\sin k(t-t_0)}{\{k(t-t_0)\}^2}\} \\ & - 5F_2(\vec{k},t_0)\{\frac{\sin k(t-t_0)}{k(t-t_0)} + \frac{3\cos k(t-t_0)}{(k(t-t_0))^2} - \frac{3\sin k(t-t_0)}{(k(t-t_0))^3}\}].\end{aligned} \tag{30}$$

This is the fluctuations in energy density, from which the fluctuations in temperature can be obtained as $\delta\rho/\bar{\rho} = 4\delta T/T$ for $\rho \sim T^4$. We use Eqs 19, 20 and 30 to obtain the density fluctuation in terms of transport coefficients as:

$$\begin{aligned}\Delta(\vec{k},t) =& e^{-(t-t_0)/\tau}[\Delta(\vec{k},t_0)\frac{\sin k(t-t_0)}{k(t-t_0)} + \frac{4}{k}\frac{\chi}{s}\frac{T(\vec{k},t_0)}{\bar{T}}\{k^2 - ik_l\dot{U}_l(\vec{k},t_0)\}\{\frac{\cos k(t-t_0)}{k(t-t_0)} - \frac{\sin k(t-t_0)}{k^2(t-t_0)^2}\} \\ & + \frac{40}{3}\frac{\eta}{s}\frac{\Theta(\vec{k},t_0)}{\bar{T}}\{\frac{\sin k(t-t_0)}{k(t-t_0)} + \frac{3\cos k(t-t_0)}{k^2(t-t_0)^2} - \frac{3\sin k(t-t_0)}{k^3(t-t_0)^3}\}].\end{aligned} \tag{31}$$

This equation provides the connection of the energy density in Fourier space with various transport coefficients *e.g.* thermal conductivity (χ) and viscosity (η). The fluctuation in energy density in position space can be obtained by taking Fourier transformation of Eq. 31 as,

$$\frac{\delta\rho}{\bar{\rho}}(\vec{x},t) = \int\frac{d^3k}{(2\pi)^3}\Delta(\vec{k},t)\exp{(i\vec{k}.\vec{x})} \tag{32}$$

If the initial ($t=t_0$) energy density fluctuation, gradient of velocity, viscosity to entropy ratio and temperature of the system in equilibrium are known then Eqs. 31 and 32 can be used to get fluctuations at any time, $t > t_0$. Angular correlation function for these fluctuation has been estimated in the appendix.

III. EVOLUTION OF FLUCTUATION IN A HYDRODYNAMICALLY EXPANDING QGP BACKGROUND

In the previous section we have considered the evolution of the fluctuations in a non-expanding background. However, in a realistic scenario in HICRE the system expands very fast due to its high internal pressure. Therefore, in this section we include the effects of the expansion on the spatial anisotropy through the solutions of relativistic hydrodynamics.

The space time variation of quantities such as energy density and flow velocity is governed by relativistic hydrodynamics. Therefore, we solve the equation:

$$\partial_\mu \bar{T}^{\mu\nu} = 0 \tag{33}$$

with the assumption that net baryon number density (n_B) is zero at the central rapidity region, hence we need not consider the equation $\partial_\mu(n_B u^\mu) = 0$. We also assume boost invariance [23] along the longitudinal direction and solve the Eq. 33 numerically for initial condition taken from optical Glauber at the highest RHIC energy ($\sqrt{s_{NN}} = 200$ GeV) for Au+Au collision. The hydrodynamic solutions [24] for flow velocity and temperature ($\bar{T} = [30\bar{\rho}/(g\pi^2)]^{1/4}$) have been used to study the space time evolution of the fluctuations in an expanding background.

The evolution of $f(x,p) = f_0(x,p) + \delta f(x,p)$ with $\delta f = f_0\psi$ is governed by the equation, $p^\mu \partial_\mu f = (p\cdot u)C[f]$ which reduces to the following for an expanding system in the relaxation time approximation:

$$\left(\frac{\partial}{\partial t} + \frac{\vec{p}}{p^0}\cdot\frac{\partial}{\partial \vec{x}} + \frac{(p^0 u_0 - \vec{p}\cdot\vec{u})}{p^0 \tau_R(x)}\right)\delta f(x,p) = -\left(\frac{\partial}{\partial t} + \frac{\vec{p}}{p^0}\cdot\frac{\partial}{\partial \vec{x}}\right) f_0(x,p) \tag{34}$$

The solution of Eq. 34 is given by [22]:

$$\delta f(x,p) = D(t,t_0)\left[\delta f_{in}(p, \vec{x} - \frac{\vec{p}}{p^0}(t-t_0)) + \int_{t_0}^{t} B(\vec{x} - \frac{\vec{p}}{p^0}(t-t'), t')\, D(t_0,t')dt'\right] \tag{35}$$

where

$$D(t_2,t_1) = \exp\left[-\int_{t_1}^{t_2} dt' A(p,\, \vec{x} - \frac{\vec{p}}{p^0}(t'-t_0),\, t')\right] \tag{36}$$

with

$$A(p,\vec{x},t) = \frac{p^0 u_0(x) - \vec{p}\cdot\vec{u}(x)}{p^0 \tau_R(x)} \tag{37}$$

and

$$B(\vec{x},t) = -\left(\frac{\partial}{\partial t} + \frac{\vec{p}}{p^0}\cdot\frac{\partial}{\partial \vec{x}}\right) f_0(x,p) \tag{38}$$

For

$$f_0(x,p) = f_{eq} = \frac{1}{e^{\beta(x)(u^\mu p_\mu) - 1}} \tag{39}$$

The expression for B reduces to:

$$B(\vec{x},t) = -f_{eq}(1+f_{eq})\frac{p^\mu}{p^0}\partial_\mu\left[\beta(x)u^\mu p_\mu\right] \tag{40}$$

We took the relaxation time as $\tau_R^{-1}(x) = 1.1\alpha_s T(x)$ [21] (we have taken constant value of $\alpha_s = 0.2$ here), $\beta = 1/T(x)$, $u^\mu(x) = (\gamma, \gamma\,\vec{v})$ is the four velocity of the fluid and $\gamma(x) = u^0(x) = (1 - v(x)^2)^{-1/2}$. The space-time variation of the temperature and velocity fields are taken from the solution of the relativistic hydrodynamic equations. To simulate different types of initial spatial anisotropy we choose,

$$\delta f(p,\vec{x},t_0) = A_0 \exp\left[-r(1 + a_n \cos n\vartheta)\right] \tag{41}$$

We have taken $n = 2,3,4,5,...$ to simulate different initial anisotropy. For numerical results A_0 is set as unity.



FIG. 1: Evolution of the fluctuation in energy density with r coordinate at different t for a non-expanding QGP background.

IV. RESULTS

We display the spatial variation of the fluctuation in Fig. 1 at different times for a non-expanding background. We substitute eq. 22 in eq. 23 with Ψ_{in} as a Gaussian in space at the initial time and evaluate the evolution of the fluctuation. For the sake of illustration we take $T = 400$ MeV resulting in $\tau \sim 0.98$ fm/c. The results indicate a rapid dissipation and displacement of the peak of the fluctuations with increase in time. Now we present the results for a realistic scenario where the background QGP is expanding hydrodynamically as described in the previous section. Taking the value of the temperature dependent relaxation time relevant for QCD plasma [21] the evolution of initial spatial anisotropies introduced through Ψ_{in} (or δf) have been studied. It is to be noted that due to expansion the temperature decreases and hence the relaxation time increases which slows down the dissipation. Therefore, the dissipation of the perturbation gets slower with the expansion of the system. In Fig. 2 the evolution of the initial elliptic spatial anisotropy (upper panel), realized by taking $n = 2$ in Eq. 41 is depicted. The initial thermalization time is taken as $\tau_0 = 0.6$ fm/c. The evolution is studied upto $\tau = 4$ fm/c when the temperature is ~ 200 MeV. For $n = 2$ the anisotropy has an elliptic shape having stronger gradient along x-axis resulting in faster expansion along x compared to y axis. Therefore, the propagation of the perturbation generates a pattern similar to the one generated in water waves by the impact of a stick on the still water surface. This kind of pattern is clearly observed in the lower panel of Fig. 2. It is also important to note that the two oppositely propagating perturbations are strongly correlated which may have interesting observation effects. Moreover, if the perturbation is created near the boundary of the system then the wave propagating outward will dissipate less than the one which will move toward the inner matter. The results indicate a rapid dissipation of the peak. The peak has been reduced by almost 70% at a time $\tau = 4$ fm/c. The displacement of the peak is determined by the sound horizon i.e. the distance travelled by the sound wave: $\int_{\tau_0}^{\tau} d\tau c_s(\tau) d\tau$. We have taken the sound velocity, $c_s = 1/\sqrt{3}$, independent of τ for the expanding QGP background. The spatial anisotropic structure of the system formed in HICRE can be understood with the help of Fourier analysis in terms of its various coefficients. Work on the space-time evolution of the angular power spectrum for more realistic initial condition for the hydrodynamical solution derived from Glauber Monte-Carlo techniques is under progress [25].

In Figs. 3 - 5 the evolutions of the spatial anisotropy with different initial geometry i.e. triangular, quadrangular and pentagonal for $n = 3, 4$ and 5 respectively have been depicted. It is observed that the spatial anisotropies dissipate fast. The propagation of these anisotropies are primarily dictated by the velocity of sound in the QGP background. The propagating waves for the perturbation take shape analogous to water waves created on the calm surface if perturbed initially with similar geometric shape. Theoretical analysis of the angular power spectrum of the anisotropies will shed light on the signatures of the early stage of the evolving matter.



FIG. 2: Evolution of the spatial anisotropy with initial elliptic geometry at time $\tau_0 = 0.6$ fm/c (upper panel). The lower panel shows the geometry after a time 4 fm/c has elapsed. Hydrodynamic expansion of the QGP background has been taken into account.

V. SUMMARY AND DISCUSSION

In the present work we have studied the time evolution of spatial anisotropies for a system formed in HICRE. The background of the spatial fluctuations has been assumed as a thermalized expanding QGP. The expansion of the background has been dealt with the (2+1) dimensional relativistic hydrodynamical model. The anisotropy has been treated within the framework of BTE in the QGP background. The evolution of initial spatial anisotropies with different geometry have been studied and it is found that the spatial anisotropies dissipate during its propagation. Theoretical analysis of these anisotropies will help in understanding the early stage of the matter. We have provided an explicit relations between the fluctuations and transport coefficients. Angular power spectrum of the anisotropies for non-expanding background has been estimated in the appendix. A detail analysis of this power spectrum with hydrodynamically expanding background is underway. This will provide detail information on the initial anisotropies.

Acknowledgement: G. S. acknowledges the support from Department of Atomic Energy, Govt. of India for this work. We would like to thank Mr Sushant Kumar Singh for providing inputs from relativistic hydrodynamical code.



FIG. 3: Same as Fig. 2 but the spatial anisotropy has a triangular geometry at the initial time $\tau_0 = 0.6$ fm/c (upper panel). The lower panel shows the geometry after a time 4 fm/c has elapsed.

VI. APPENDIX: CORRELATIONS

In this appendix, we evaluate the correlation, $\langle\Delta(\hat{n_1})\Delta(\hat{n_2})\rangle$. The fluctuations, $\Delta(\vec{x},\hat{n},t)$ can be written as:

$$\Delta(\vec{x},\hat{n},t) = \sum_{l,m} a_{lm}(\vec{x},t) Y_{lm}(\hat{n}), \tag{42}$$

with $a_{lm} = (-i)^l 4\pi \int d^3\hat{n}\, Y^*_{lm}(\hat{n})\Delta(\vec{x},\hat{n},t)$ and $\langle a_{lm}.a^*_{l'm'}\rangle = C_l(\vec{x},t)\delta_{ll'}\delta_{mm'}$ Using Eq. 42 we can find the correlations in the following way,

$$\begin{aligned}
\langle\Delta(\vec{x},\hat{n}_1,t)\Delta(\vec{x},\hat{n}_2,t)\rangle &= \sum_{l,m,l',m'} \langle a_{lm}.a^*_{l'm'}\rangle Y_{lm}(\hat{n}_1) Y^*_{l'm'}(\hat{n}_2), \\
&= \sum_{l,m,l',m'} C_l(\vec{x},t)\delta_{ll'}\delta_{mm'} Y_{lm}(\hat{n}_1) Y^*_{l'm'}(\hat{n}_2), \\
&= \sum_{l,m} C_l(\vec{x},t) Y_{lm}(\hat{n}_1) Y^*_{lm}(\hat{n}_2),
\end{aligned} \tag{43}$$

giving $\langle\Delta(\vec{x},\hat{n}_1,t)\Delta(\vec{x},\hat{n}_2,t)\rangle = \frac{1}{4\pi}\sum_l (2l+1) C_l(\vec{x},t) P_l(\hat{n}_1\cdot\hat{n}_2)$, since, $P_l(\hat{n}_1\cdot\hat{n}_2) = \frac{4\pi}{2l+1}\sum_{m=-l}^{l} Y_{lm}(\hat{n}_1) Y^*_{lm}(\hat{n}_2)$. Eq. 43 defines the correlations of fluctuations observed from two different directions in terms of co-efficients C_ls. C_l's are the angular power spectrum which contains the information of the anisotropies. Work is under progress to estimate these coefficients by solving the



FIG. 4: Same as Fig. 2 but the spatial anisotropy has a quadrangular geometry at the initial time $\tau_0 = 0.6$ fm/c (upper panel). The lower panel shows the geometry after a time 4 fm/c has elapsed.

hydrodynamical equations with the initial conditions taken from Glauber Monte-Carlo method. Similarly, one can define these co-efficients corresponding to k-space presentation of fluctuations. Time evolution of these co-efficients can be obtained from evolution of energy density fluctuation, $\Delta(\vec{k}, \hat{n}, t)$ given by Eq. 31. Therefore, we have

$$\langle \Delta(\vec{x}, \hat{n}_1, t)\Delta(\vec{x}, \hat{n}_2, t)\rangle = \int \frac{d^3k}{(2\pi)^3}\frac{d^3k'}{(2\pi)^3}\; e^{i(\vec{k}-\vec{k'})\cdot\vec{x}}\; \langle \Delta(\vec{k}, \hat{n}_1, t)\Delta(\vec{k'}, \hat{n}_2, t)\rangle \tag{44}$$

Now,

$$\begin{aligned}\langle \Delta(\vec{k}, \hat{n}_1, t)\Delta(\vec{k'}, \hat{n}_2, t)\rangle =& \langle \Delta(\vec{x}, \hat{n}_1, t_0)\Delta(\vec{x}, \hat{n}_2, t_0)\rangle L(\vec{k}, t_0; \vec{k'}, t_0)\\ &+ \langle \Theta(\vec{x}, \hat{n}_1, t_0)\Theta(\vec{x}, \hat{n}_2, t_0)\rangle M(\vec{k}, t_0; \vec{k'}, t_0)\\ &+ \langle \Delta(\vec{x}, \hat{n}_1, t_0)\Theta(\vec{x}, \hat{n}_2, t_0)\rangle N(\vec{k}, t_0; \vec{k'}, t_0),\end{aligned} \tag{45}$$



FIG. 5: Same as Fig. 2 but the spatial anisotropy has a pentagonal geometry at the initial time $\tau_0 = 0.6$ fm/c (upper panel). The lower panel shows the geometry after a time 4 fm/c has elapsed.

where,

$$\begin{aligned}
L(\vec{k},t,\vec{k}',t_0) =& e^{-\frac{2(t-t_0)}{\tau}}\{\frac{\sin k(t-t_0)}{k(t-t_0)}\}\{\frac{\sin k'(t-t_0)}{k'(t-t_0)}\},\\
M(\vec{k},t,\vec{k}',t_0) =& e^{-\frac{2(t-t_0)}{\tau}}\left(\frac{40}{3}\frac{\eta}{s\bar{T}}\right)^2\{\frac{\sin k(t-t_0)}{k(t-t_0)}+\frac{3\cos k(t-t_0)}{(k(t-t_0))^2}-\frac{3\sin k(t-t_0)}{(k(t-t_0))^3}\}\\
&\times\{\frac{\sin k'(t-t_0)}{k'(t-t_0)}+\frac{3\cos k'(t-t_0)}{(k'(t-t_0))^2}-\frac{3\sin k'(t-t_0)}{(k'(t-t_0))^3}\} \qquad (46)\\
N(\vec{k},t,\vec{k}',t_0) =& e^{-\frac{2(t-t_0)}{\tau}}\frac{40}{3}(\frac{\eta}{s\bar{T}})\{\frac{\sin k(t-t_0)}{k(t-t_0)}\}\\
&\times\{\frac{\sin k'(t-t_0)}{k'(t-t_0)}+\frac{3\cos k'(t-t_0)}{(k'(t-t_0))^2}-\frac{3\sin k'(t-t_0)}{(k'(t-t_0))^3}\}
\end{aligned}$$

For two functions Δ and Θ, defining the correlation as: $\langle\Delta(\vec{k},\hat{n}_1,t)\Theta(\vec{k}',\hat{n}_2,t)\rangle = (2\pi)^3\delta(\vec{k}-\vec{k}')\delta_{\Delta\Theta}\langle\Delta(k,\hat{n}_1,t)\Theta(k,\hat{n}_2,t)\rangle$ and $\Delta(\vec{k},\hat{n},t)=\sum_{l,m}a^{\Delta}_{lm}(\vec{k},t)Y_{lm}(\hat{n})$, $\langle {a^{\Delta}}_{lm}.{a^{*}}^{\Theta}_{l'm'}\rangle = C^{\Delta\Theta}_l(\vec{k},t)\delta_{ll'}\delta_{mm'}$, we get

$$\begin{aligned}
\langle\Delta(\vec{k},\hat{n}_1,t)\Delta(\vec{k}',\hat{n}_2,t)\rangle =& (2\pi)^3\delta(\vec{k}-\vec{k}')\sum_l\frac{2l+1}{4\pi}P_l(\hat{n}_1\cdot\hat{n}_2)\\
&\times\{C^{\Delta\Delta}_l(k,t_0)L(k,t,k,t_0)+C^{\Theta\Theta}_l(k,t_0)M(k,t,k,t_0)\}. \qquad (47)
\end{aligned}$$

Using $\langle\Delta(\vec{k},\hat{n}_1,t)\Delta(\vec{k'},\hat{n}_2,t)\rangle = (2\pi)^3\delta(\vec{k}-\vec{k'})\sum_l \frac{2l+1}{4\pi}C_l^{\Delta\Delta}(k,t)P_l(\hat{n}_1\cdot\hat{n}_2)$, in Eq. 47 we get,

$$C_l^{\Delta\Delta}(\vec{k},t) = C_l^{\Delta\Delta}(k,t_0)L(k,t,k,t_0) + C_l^{\Theta\Theta}(k,t_0)M(k,t,k,t_0). \tag{48}$$

Using Eq. 43, 48 in Eq 44, we get

$$C_l^{\Delta\Delta}(\vec{x},t) = \int d^3k\{C_l^{\Delta\Delta}(k,t_0)L(k,t,k,t_0) + C_l^{\Theta\Theta}(k,t_0)M(k,t,k,t_0)\} \tag{49}$$

This provides the correlations at different angular scales at time t for a given correlations at initial time, t_0 $(t > t_0)$.

---

[1] R. Durrer, The Cosmic Microwave Background, Cambridge University Press, 2008.
[2] P. Staig and E. Shuryak, Phys. Rev. C **84**, 034908; Phys. Rev. C **84**, 044912 (2011).
[3] A. P. Mishra, R. K. Mohapatra, P. S. Saumia and A. M. Srivastava, Phys. Rev. C **81**, 034903 (2010).
[4] G.-Y. Qin and B. Mueller, Phys. Rev. C **85**, 061901(R) (2012).
[5] S. R. de Groot, W. A. van Leeuwen and Ch. G. van Weert, Relativistic Kinetic Theory, North Holland Publishing Co., 1980.
[6] M. A. Stephanov, K. Rajagopal and E. V. Shuryak, Phys. Rev. Lett. **81**, 4816 (1998).
[7] M. A. Stephanov, K. Rajagopal and E. V. Shuryak, Phys. Rev. D **60**, 114028 (1999).
[8] M. Asakawa, U. Heinz and B. Muller, Phys. Rev. Lett. **85**, 2072 (2000).
[9] S. Jeon and V. Koch, Phys. Rev. Lett. **83**, 5435 (1999).
[10] S. Jeon and V. Koch, Phys. Rev. Lett. **85**, 2076 (2000).
[11] M. A. Stephanov, Phys. Rev. D **81**, 054012 (2010).
[12] S. Mrowczynski, Phys. Rev. C **57**, 1518 (1998).
[13] A. Rafiei and K. Javidan, Phys. Rev. C **94**, 034904 (2016).
[14] E. M. Lifshitz and L. P. Pitaevskii, Physical Kinetics (Elsevier, New York, 1987).
[15] S. Weinberg, Gravitation and Cosmology (John Wiley & Sons, New York, 1972)
[16] E. Flowers and N. Itoh, Astrophys. J. **230**, 847 (1979).
[17] C. -P. Ma and E. Bertschinger, Astro. Phys. J. **455**, 7 (1995).
[18] W. Florkowski, R. Ryblewski, M. Strickland, Phys. Rev. C **88**, 024903 (2013).
[19] W. Florkowski, R. Ryblewski, M. Strickland, Phys. Rev. D **86**, 085023 (2012).
[20] Y. Hatta, M. Martinez and B. -W. Xiao, Phys. Rev. D **91**. 085024 (2015).
[21] M. H. Thoma, Phys. Rev. D **49**, 451 (1994).
[22] H. Levine, Partial Differential Equations, American Mathematical Society (1997).
[23] J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).
[24] S. K. Singh *et al.* under preparation.
[25] G. Sarwar and J. Alam (to be published).